

## SPECIAL ARTICLE

### UNCONVENTIONAL MEDICINE IN THE UNITED STATES

#### Prevalence, Costs, and Patterns of Use

DAVID M. EISENBERG, M.D., RONALD C. KESSLER, PH.D., CINDY FOSTER, M.P.H.,  
FRANCES E. NORLOCK, M.P.H., DAVID R. CALKINS, M.D., M.P.P., AND THOMAS L. DELBANCO, M.D.

**Abstract Background.** Many people use unconventional therapies for health problems, but the extent of this use and the costs are not known. We conducted a national survey to determine the prevalence, costs, and patterns of use of unconventional therapies, such as acupuncture and chiropractic.

**Methods.** We limited the therapies studied to 16 commonly used interventions neither taught widely in U.S. medical schools nor generally available in U.S. hospitals. We completed telephone interviews with 1539 adults (response rate, 67 percent) in a national sample of adults 18 years of age or older in 1990. We asked respondents to report any serious or bothersome medical conditions and details of their use of conventional medical services; we then inquired about their use of unconventional therapy.

**Results.** One in three respondents (34 percent) reported using at least one unconventional therapy in the past year, and a third of these saw providers for unconventional therapy. The latter group had made an average of 19 visits to such providers during the preceding year, with an average charge per visit of \$27.60. The frequency of use of unconventional therapy varied somewhat among sociodemographic groups, with the highest use reported by

nonblack persons from 25 to 49 years of age who had relatively more education and higher incomes. The majority used unconventional therapy for chronic, as opposed to life-threatening, medical conditions. Among those who used unconventional therapy for serious medical conditions, the vast majority (83 percent) also sought treatment for the same condition from a medical doctor; however, 72 percent of the respondents who used unconventional therapy did not inform their medical doctor that they had done so. Extrapolation to the U.S. population suggests that in 1990 Americans made an estimated 425 million visits to providers of unconventional therapy. This number exceeds the number of visits to all U.S. primary care physicians (388 million). Expenditures associated with use of unconventional therapy in 1990 amounted to approximately \$13.7 billion, three quarters of which (\$10.3 billion) was paid out of pocket. This figure is comparable to the \$12.8 billion spent out of pocket annually for all hospitalizations in the United States.

**Conclusions.** The frequency of use of unconventional therapy in the United States is far higher than previously reported. Medical doctors should ask about their patients' use of unconventional therapy whenever they obtain a medical history. (N Engl J Med 1993;328:246-52.)

UNCONVENTIONAL, alternative, or unorthodox therapies are difficult to define, because they encompass a broad spectrum of practices and beliefs. As Murray and Rubel have written, "Many are well known, others are exotic or mysterious, and some are dangerous."<sup>1</sup> From a sociological standpoint, unconventional therapy refers to medical practices that are not in conformity with the standards of the medical community.<sup>2</sup> Here we define unconventional therapies as medical interventions not taught widely at U.S. medical schools or generally available at U.S. hospitals. Examples include acupuncture, chiropractic, and massage therapy.

Studies based on samples in limited geographic areas suggest that the use of unconventional therapy is widespread.<sup>3-5</sup> In particular, unconventional therapies are frequently used by patients with cancer,<sup>6-11</sup> arthritis,<sup>11-13</sup> chronic back pain,<sup>3,14</sup> the acquired immunodeficiency syndrome,<sup>15</sup> gastrointestinal problems,<sup>16,17</sup>

chronic renal failure,<sup>18</sup> and eating disorders.<sup>19</sup> Little is known, however, about the overall prevalence, cost, and patterns of use of unconventional therapy in the United States.<sup>1,20</sup>

To improve our understanding of the use of unconventional therapy, we conducted a national telephone survey focusing on 16 interventions found, on the basis of pilot research, to be representative of unconventional therapies used commonly in the United States. Our study focused on the following questions: What is the extent of use of unconventional therapy in the United States? How much is spent annually on these therapies, including out-of-pocket and third-party payments? What sociodemographic factors distinguish users of unconventional therapy from nonusers? For what medical conditions do people most commonly use unconventional therapy? And to what extent are medical doctors responsible for or informed about the use of unconventional therapy by their patients?

#### METHODS

##### Sample

We conducted our survey by telephone between January 18 and March 7, 1991. The sample was selected by means of random-digit dialing.<sup>21</sup> We limited eligibility to English-speaking persons, 18 years of age or older, in whom cognitive or physical impairment did not prevent the completion of the interview. We designed the survey with a target sample of 1500. Assuming an estimated prevalence of use of unconventional therapy between 10 and 50 percent, we calcu-

From the Division of General Medicine and Primary Care, Department of Medicine, Beth Israel Hospital and Harvard Medical School, Boston (D.M.E., T.L.D.); the Institute for Social Research, University of Michigan, Ann Arbor (R.C.K., C.F.); the Division of General Medicine, Department of Medicine, New England Deaconess Hospital and Harvard Medical School, Boston (D.R.C.); and Chicago College for Osteopathic Medicine, Chicago (F.E.N.). Address reprint requests to Dr. Eisenberg at the Division of General Medicine and Primary Care, Beth Israel Hospital, 330 Brookline Ave., Boston, MA 02215.

Supported by a contract with the John E. Fetzer Institute and a grant from the Nathan Cummings Foundation.

lated that 1500 interviews should produce estimated prevalence rates with 95 percent confidence intervals of 2 to 3 percent.

Of the initial sample of 5158 telephone numbers, 38 percent were nonworking, and 13 percent were not assigned to households. We declared 221 respondents ineligible because they did not speak English (97), because of cognitive or physical incapacity (96), or because they were temporarily unavailable (28). Among the remaining 2295 eligible respondents, 1539 completed the interview, 653 declined to participate (81 of them before we could establish eligibility), and 103 began the interview but stopped before completing all questions. These figures correspond to a 67 percent overall response rate among eligible respondents. Only one respondent per household was eligible to be interviewed. This person was selected by computer randomization from the list of household members given by the first household member contacted. Persons with responses substantially different from the remainder of the sample (for example, those with frequent visits to a provider of unconventional therapy at no cost) were contacted again by a supervisor for verification or clarification of their responses. Since we asked respondents about the use of unconventional therapy during the 12 months before the interview, we considered the results representative of 1990.

### The Interview

We described the interview to the respondents as a survey by investigators from Harvard Medical School that was designed to assess the health care practices of Americans. We made no mention of unconventional therapy while recruiting the respondents. The interviews, which averaged 25 minutes in length, began with questions on the respondents' health, health worries, days in bed at home or in the hospital, and indicators of functional impairment caused by health problems. We then asked the respondents about their interactions with medical doctors during the past 12 months. A "medical doctor" was defined early in the interview as "a medical doctor (M.D.) or an osteopath (D.O.), not a chiropractor or other nonmedical doctor." Throughout the remainder of the interview we used the term "medical doctor." We use the same term in this report when referring to a respondent's provider of conventional medical care.

We next assessed the respondents' medical problems. The interviewers stated: "Now I'm going to read a list of conditions. Please tell me if you have had any of these conditions in the past 12 months." The interviewers then asked about 24 medical conditions and offered a follow-up question, "What other important conditions did you have?" The 24 conditions included common symptoms (such as back problems, digestive problems, dizziness, headache, and allergies), as well as specific diagnoses (such as high blood pressure, diabetes, and cancer). Only 8 percent of the respondents reported conditions not included in our list.

The respondents were then asked to identify the three (or fewer) "most bothersome or serious" health problems from the list they had just provided. These health problems are referred to here as "principal medical conditions." We asked the respondents whether they had seen their medical doctor for each principal medical condition during the past 12 months and what their perceptions of these interactions had been.

At this point we asked respondents about their use of unconventional therapy. The interviewers' text read as follows: "Now I'd like to ask you about your use of some other kinds of therapies and treatments." The respondents were asked whether they had ever used 1 or more of 16 unconventional therapies for their principal medical conditions and, if so, whether they had used any of them in the past 12 months. For example, a respondent listing back problems as a principal medical condition was asked whether he or she had used any of the 16 unconventional therapies for this problem during the past 12 months. The respondents were next asked whether they had used "any other therapy not generally provided by most clinics and hospitals." Only a small number (1 percent of the sample) reported using any other unconventional therapy.

Some of the unconventional therapies studied warrant further clarification. For example, "massage therapy" or "relaxation therapy" may mean different things to different people. A respondent who used "massage therapy" for a specific principal medical condi-

tion was asked, "Could you tell me more about your use of massage therapy? For example, what technique do you use?" Similar questions followed a report of the use of herbal therapy, spiritual or religious healing by others, commercial weight-loss programs, lifestyle diets, energy healing, folk remedies, and megavitamin therapy. (With regard to megavitamin use, the interviewers specifically stated that "megavitamin therapy does not include taking a daily vitamin.") Given that some unconventional therapies, such as massage, were available in more than 100 reported varieties, further subclassification of each of the 16 unconventional therapies is beyond the scope of this paper.

When respondents reported the use of unconventional therapy during the past 12 months, we asked whether a "professional" was involved. Specifically, the interviewers described a professional as "someone who provides care or gives advice and is paid for his or her services." Such persons are referred to in this report as "providers of unconventional therapy." Some forms of unconventional therapy typically involve a provider (for example, a chiropractor or acupuncturist), whereas others do not (for example, lifestyle diet or self-help groups). In addition, some users of unconventional therapy may visit a provider less often than once a year but may continue to use the prescribed unconventional therapy. In order to learn more about these variations, we asked about lifetime visits and recent visits (in the past 12 months) to providers of unconventional therapy. We did not ask whether the respondents' providers of unconventional therapy were medical doctors. We asked about total charges for visits to providers of unconventional therapy during the past 12 months and whether insurance paid any of these charges. The interviewers asked whether the respondents had discussed the use of each unconventional therapy with their medical doctors and collected data on the respondents' demographic characteristics. We also asked questions pertaining to the respondents' perceptions of the efficacy of unconventional therapies and the quality of their interactions with the providers. We shall report these data separately.

Pilot research suggested that, in addition to the 16 unconventional therapies we studied, prayer and exercise are commonly used in the United States for purposes related to health. We therefore asked respondents whether they had used prayer or exercise as a medical "therapy or treatment" in the past year. No additional questions were asked about these two activities. In the case of prayer, we thought such questions would be inappropriate. As for exercise, we thought the term was too vague and the practice sufficiently ubiquitous to preclude the gathering of useful data. Unless they are explicitly mentioned, all analyses described here excluded prayer and exercise.

### Statistical Analysis

We weighted the data to adjust for variations among households in the number of telephones and number of household members eligible to participate in the survey. We also weighted the data to match our sample to the distribution of the U.S. population, as reported in the U.S. Census, for age, sex, and education. Using the Taylor series approximation method for calculating standard errors, we carried out tests of significance appropriate for weighted data using the SUDAAN software system.<sup>22</sup> We based our extrapolations of the estimates from the survey to the total U.S. household population (which, according to the U.S. Census Bureau, excludes homeless and institutionalized persons) on preliminary figures from the 1990 U.S. Census, which reports the total U.S. population as 242 million, with 74 percent (approximately 180 million persons) made up of adults 18 years of age or older living in U.S. households.

## RESULTS

### Characteristics of the Respondents and Generalizability of the Sample

The characteristics of the subjects we interviewed are shown in Table 1. The sociodemographic characteristics of the survey population were similar to those

in the 1989 U.S. National Health Interview Survey<sup>23</sup> with respect to age, sex, race, education, marital status, and region of the country.

Table 2 summarizes the use of unconventional therapy in the 12 months before the survey for all 1539 respondents interviewed. Excluding exercise and prayer, one in three respondents (34 percent) used at least one unconventional therapy in 1990. Nearly two thirds (64 percent) of those who used unconventional therapy did so without visiting a provider of unconventional therapy during the 12 months before the interview, whereas the one third who did see a provider made an average of 19 visits.

The use of unconventional therapy was not confined to any narrow segment of U.S. society. The rates of use ranged from 23 to 53 percent in all sociodemographic groups we considered. There were no significant differences according to sex or insurance status

Table 1. Characteristics of the 1539 Subjects Interviewed.

CHARACTERISTIC	No. (%)*
<b>Sex</b>	
Female	732 (48)
Male	807 (52)
<b>Age (yr)</b>	
18-24	247 (16)
25-34	354 (23)
35-49	409 (27)
≥50	529 (34)
<b>Race or ethnic group</b>	
White	1264 (82)
Black	140 (9)
Hispanic	95 (6)
Asian	16 (1)
Other	24 (2)
<b>Education</b>	
<High school	370 (24)
High school	543 (35)
College or trade school	345 (22)
Graduate school	281 (18)
<b>Annual income</b>	
<\$20,000	462 (30)
\$20,000-34,999	517 (34)
\$35,000-49,999	289 (19)
≥50,000	271 (18)
<b>Size of community†</b>	
Big city	170 (11)
Small city	433 (28)
Big suburb	124 (8)
Small suburb	186 (12)
Town	274 (18)
Rural area	352 (23)
<b>Region</b>	
East	340 (22)
South	498 (32)
Midwest	401 (26)
West	300 (19)
<b>No. of principal medical conditions reported</b>	
None	261 (17)
1	290 (19)
2	242 (16)
≥3	746 (48)
At least 1	1279 (83)

\*Because of rounding, percentages do not always total 100.

†Defined in terms of population: big city, ≥1 million; small city, <1 million; big suburb, suburb of a city with population ≥1 million; and small suburb, suburb of a city with population <1 million.

Table 2. Prevalence and Frequency of Use of Unconventional Therapy among 1539 Adult Respondents in 1990.

TYPE OF THERAPY	USED IN PAST 12 Mo (%)*	SAW A PROVIDER (%)*	MEAN NO. OF VISITS PER USER IN PAST 12 Mo
Relaxation techniques	13	9	19
Chiropractic	10	70	13
Massage	7	41	15
Imagery	4	15	14
Spiritual healing	4	9	14
Commercial weight-loss programs	4	24	23
Lifestyle diets (e.g., macrobiotics)	4	13	8
Herbal medicine	3	10	8
Megavitamin therapy	2	12	13
Self-help groups	2	38	21
Energy healing	1	32	8
Biofeedback	1	21	6
Hypnosis	1	52	3
Homeopathy	1	32	6
Acupuncture	<1	91	38
Folk remedies	<1	0	0
Exercise‡	26	—	—
Prayer‡	25	—	—
≥1 Unconventional therapy‡	34	36	19
95% Confidence interval	31-37	31-41	14-24

\*Percentages are of those who used that type of unconventional therapy.

†Respondents who used exercise or prayer were not asked for details about this use.

‡Excluding exercise and prayer.

and only small variations according to the size of the community. The use of unconventional therapy was significantly more common among people 25 to 49 years of age (38 percent) than among those who were younger (33 percent) or older (28 percent) ( $P<0.05$  for both comparisons). The use of unconventional therapy was significantly less common among blacks (23 percent) than among members of other racial groups (35 percent;  $P<0.05$ ). It was significantly more common among persons with some college education (44 percent) than among those with no college education (27 percent;  $P<0.05$ ) and significantly more common among people with annual incomes above \$35,000 (39 percent) than among those with lower incomes (31 percent;  $P<0.05$ ). Use was also significantly more common among those living in the West (44 percent) than among those living in the rest of the country (31 percent;  $P<0.05$ ).

#### Patterns of Use

The vast majority of respondents (83 percent) reported one or more principal medical conditions in 1990. More than half (58 percent) of respondents with at least one principal medical condition saw a medical doctor but not a provider of unconventional therapy in 1990; 3 percent saw only a provider of unconventional therapy; 7 percent saw both a medical doctor and a provider of unconventional therapy; and 33 percent saw neither for at least one principal medical condition (Fig. 1).

Table 3 summarizes the rates of use of unconventional therapy for the 10 most common principal medical conditions. On average, 1 in 4 respondents




Figure 1. Percentage of Respondents Reporting at Least One Principal Medical Condition Who Saw a Medical Doctor or Provider of Unconventional Therapy in 1990.

Eighty-three percent (1279) of the 1539 respondents reported one or more principal medical conditions. "Provider" denotes a provider of unconventional therapy.

(25 percent) used unconventional therapy and 1 in 10 (10 percent) went to a provider of unconventional therapy for a principal medical condition in 1990. Among all the conditions studied, the frequency of use of unconventional therapy was highest for back problems (36 percent), anxiety (28 percent), headaches (27 percent), chronic pain (26 percent), and cancer or tumors (24 percent). Consistent with the prevalence of cancer in the general population, the use of unconventional therapy for cancer accounted for less than 3 percent of all use. Relaxation techniques, chiropractic, and massage were the unconventional therapies used most often in 1990.

Among respondents who reported a principal medical condition and used unconventional therapy for that condition, only 4 percent saw a provider of unconventional therapy without also seeing a medical doctor. No respondent saw a provider of unconventional therapy, but not a medical doctor, for the treatment of cancer, diabetes, lung problems, skin problems, high blood pressure, urinary tract problems, or dental problems.

To clarify further the patterns of use of unconventional therapy, we defined and investigated two predominant patterns of care for each principal medical condition: the conventional pattern (in which the respondent sought the services of a medical doctor but did not use unconventional therapies) and the unconventional pattern (the respondent used unconventional therapy with or without seeing a medical doctor). The unconventional pattern was more common than the conventional pattern for 5 of the 10 most frequently cited principal

medical conditions: back problems, insomnia, headache, anxiety, and depression.

Among respondents who saw a medical doctor for at least one principal medical condition, more than one in four (28 percent) also used unconventional therapy, and one in nine (11 percent) saw a provider of unconventional therapy for the same condition during the 12 months before the survey (Fig. 2). The rates of use of unconventional therapy among those who consulted a medical doctor varied from condition to condition. The probability that an individual patient who saw a medical doctor also used unconventional therapy in 1990 was higher than one in three for patients with anxiety (45 percent), obesity (41 percent), back problems (36 percent), depression (35 percent), or chronic pain (34 percent). Respondents who consulted a medical doctor used unconventional therapy least often for diabetes (2 percent), dermatologic problems (7 percent), urinary problems (10 percent), gynecologic problems (11 percent), dental problems (11 percent), pulmonary problems (11 percent), and high blood pressure (12 percent).

Almost 9 of 10 respondents (89 percent) who saw a provider of unconventional therapy in 1990 did so without the recommendation of their medical doctor. In more than 7 of 10 instances (72 percent), users of unconventional therapy did not inform their medical doctor of their use of the therapy. Medical doctors were most likely to be informed about the use of homeopathy (73 percent), megavitamin therapy (72 percent), and self-help groups (61 percent) and least likely to be informed about folk remedies (11 percent), religious or spiritual healing by others (17 percent), or imagery (19 percent).

As shown in Figure 1, respondents with one or more principal medical conditions were far more likely to see a medical doctor (65 percent) than a provider of unconventional therapy (10 percent). We observed a similar pattern favoring conventional medical care among the group of respondents who reported prin-

Table 3. Use of Unconventional Therapy for the 10 Most Frequently Reported Principal Medical Conditions.

CONDITION	PERCENT REPORTING CONDITION	USED UNCONVENTIONAL THERAPY IN PAST 12 Mo*	SAW PROVIDER IN PAST 12 Mo*	THERAPIES MOST COMMONLY USED
Back problems	20	36	19	Chiropractic, massage
Allergies	16	9	3	Spiritual healing, lifestyle diet
Arthritis	16	18	7	Chiropractic, relaxation techniques
Insomnia	14	20	4	Relaxation techniques, imagery
Sprains or strains	13	22	10	Massage, relaxation techniques
Headache	13	27	6	Relaxation techniques, chiropractic
High blood pressure	11	11	3	Relaxation techniques, homeopathy
Digestive problems	10	13	4	Relaxation techniques, megavitamins
Anxiety	10	28	6	Relaxation techniques, imagery
Depression	8	20	7	Relaxation techniques, self-help groups
10 most common	73	25	10	Relaxation techniques, chiropractic, massage

\*Percentages are of those who reported the condition. "Provider" denotes a provider of unconventional therapy.

principal medical conditions and who used unconventional therapy for these conditions. Such persons were far more likely to have sought treatment from a conventional medical doctor (83 percent) during the past 12 months than they were to have seen a provider of unconventional therapy (36 percent). However, as mentioned earlier, the majority of users of unconventional therapy did not inform their medical doctors of their use of unconventional therapy. As a result, nearly half (47 percent) of respondents who used unconventional therapy for their principal medical condition did so without any professional supervision; that is, without either visiting a provider of unconventional therapy or discussing their unconventional therapy with their medical doctor.

#### Payment for Unconventional Therapy

Data on reimbursement for expenditures for unconventional therapy are shown in Table 4. The majority of respondents (55 percent) paid the entire cost of their visits out of pocket. Third-party payment was most common for the services of herbal therapists (83 percent), providers of biofeedback (40 percent), chiropractors (39 percent), and providers of megavitamins (30 percent).

#### National Projections of Use and Expenditures

Extrapolation to the total U.S. household population suggests that in 1990 an estimated 61 million Americans used at least 1 of the 16 unconventional therapies we studied and approximately 22 million Americans saw providers of unconventional therapy for a principal medical condition. Fourteen of the 16 unconventional therapies studied were used by an estimated million or more persons in 1990.

The estimated number of ambulatory visits to pro-

Table 4. Payment for Unconventional Therapy in the United States in 1990.

CATEGORY	PERCENT OF SERVICES
Unreimbursed by third-party payer	55
Partially reimbursed by third-party payer	31
Totally reimbursed by third-party payer	14
	PERCENT OF CHARGES*
Paid for out of pocket	70
	DOLLARS
Mean charge per visit to provider	27.60†
Mean out-of-pocket payment per visit to provider	19.39

\*Includes only charges for the services of providers of unconventional therapy.

†Charges ranged from <\$20 per visit for imagery, self-help, commercial weight-loss programs, spiritual healing, and acupuncture to >\$106 per visit for energy healing.

viders of unconventional therapy in 1990 was 425 million (95 percent confidence interval, 302 million to 548 million). This number exceeds the estimated 388 million visits in 1990 to all primary care physicians (general and family practitioners, pediatricians, and specialists in internal medicine) combined.<sup>24</sup>

National projections of expenditures for unconventional therapy are summarized in Table 5. If one assumes that charges for visits to providers of unconventional therapy were paid in full, Americans spent approximately \$11.7 billion for these services in 1990. This estimate refers only to the services of providers and does not include expenditures for drugs such as herbs or for medical equipment, devices, books, and other materials.

We investigated two additional expenditures by asking respondents about their use of commercial diet supplements (such as instant diet formulas, diet pills, and prepackaged meals) and over-the-counter megavitamins. Respondents who used these supplements reported out-of-pocket expenditures averaging \$228 per person per year for diet supplements and \$203 per person per year for megavitamins. These results yield national projections of approximately \$1.2 billion and \$0.8 billion, respectively. Adding these supplemental expenses to the projected expenditures for all visits to providers of unconventional therapy, we estimate that expenditures in 1990 amounted to \$13.7 billion.

The total projected out-of-pocket expenditure for unconventional therapy plus supplements was \$10.3 billion in 1990. This is comparable to the out-of-pocket expenditure for all hospital care in the




Figure 2. Use of Unconventional Therapy by Respondents Who Saw a Medical Doctor for a Principal Medical Condition in 1990.

The 10 most commonly reported conditions are shown in descending order of prevalence.

Table 5. National Projections of Expenditures for Unconventional Therapy in the United States in 1990.

CATEGORY OF EXPENDITURE	BILLIONS OF DOLLARS*
Services of providers of unconventional therapy	11.7
Megavitamin supplements	0.8
Commercial diet supplements	1.2
Estimated total	13.7
<b>Out-of-pocket expenditures</b>	
Unconventional therapy	10.3
All hospitalizations	12.8†
All physicians' services	23.5†

\*All amounts are estimates.

†From Levit et al.<sup>25</sup>

United States in 1990 (\$12.8 billion), and it is nearly half the amount spent out of pocket for all physicians' services in the United States (\$23.5 billion).<sup>25</sup>

### DISCUSSION

We found that unconventional medicine has an enormous presence in the U.S. health care system. An estimated one in three persons in the U.S. adult population used unconventional therapy in 1990. The estimated number of visits made in 1990 to providers of unconventional therapy was greater than the number of visits to all primary care medical doctors nationwide, and the amount spent out of pocket on unconventional therapy was comparable to the amount spent out of pocket by Americans for all hospitalizations. Roughly 1 in 4 Americans who see their medical doctors for a serious health problem may be using unconventional therapy in addition to conventional medicine for that problem, and 7 of 10 such encounters take place without patients' telling their medical doctors that they use unconventional therapy. Furthermore, use is distributed widely across all sociodemographic groups.

There are limits to the representativeness of our sample because it was confined to households with telephones. People living in households without telephones, those in shelters or on the street, and those in institutions were not sampled. In addition, we excluded non-English-speakers and persons for whom the interview would be burdensome because of physical or mental impairment. The frequency and patterns of use of unconventional therapy among these subgroups (and among children) are not known.

As regards the generalizability of the responses of the 67 percent of the respondents who completed the interview, we made two sets of comparisons with pre-existing national surveys. Our sample corresponded with the distribution of the subjects of the National Health Interview Survey<sup>23</sup> with respect to age, sex, race, social class, and other sociodemographic variables, suggesting that our sample was representative of the U.S. household population. Compared with the national Health and Nutrition Examination Survey,<sup>26</sup> however, our survey involved fewer people who re-

ported poor health on a five-point scale of health status (3 percent vs. 7 percent). This underrepresentation of respondents with poor health reflects our exclusion of those for whom the survey would have been burdensome because of their physical incapacity. Since we found that persons who reported poor health had substantially higher rates of use of unconventional therapy than those who perceived themselves to be in better health (52 percent vs. 33 percent), the study design may have resulted in an underestimate of the use of unconventional therapy.

Unconventional therapies are generally used as adjuncts to conventional therapy, rather than as replacements for it. Users of unconventional therapy were more likely to see a medical doctor than a provider of unconventional therapy, and visits to providers for serious medical conditions in the absence of contact with a medical doctor were rare. Moreover, in contrast to previous reports of research involving patients with cancer,<sup>6,7,9</sup> no respondents in this national survey who identified cancer as a principal medical problem reported seeing a provider of unconventional therapy without also seeing a medical doctor for this condition.

Although much of the literature dealing with unconventional medical practices focuses on potentially life-threatening or debilitating illness, the use of unconventional therapy was not limited to life-threatening conditions. Projections from survey data indicate that millions of Americans used unconventional therapy for each of the conditions we studied, the majority of which are not life-threatening. Put somewhat differently, it is likely that virtually all medical doctors see patients who routinely use unconventional therapies. Indeed, for medical doctors currently caring for patients with back problems, anxiety, depression, or chronic pain, the odds are greater than one in three that a patient is simultaneously using unconventional therapy for these medical problems without disclosing this fact.

Our results also suggest that the use of unconventional therapy is not limited to the person's principal medical conditions. A full third of the respondents who used unconventional therapy in 1990 did not use it for any of their principal medical conditions. From this fact we infer that a substantial amount of unconventional therapy is used for nonserious medical conditions, health promotion, or disease prevention. However, these issues were not a focus of our inquiry.

Although users of unconventional therapy are more likely to be in contact with medical doctors than with providers of unconventional therapy, fewer than 3 in 10 users of unconventional therapy mention its use to their medical doctors. Moreover, roughly half of those who use unconventional therapy for their principal medical conditions have no supervision of this treatment by either a medical doctor or a provider of unconventional therapy. Extrapolations to the U.S. population suggest that approximately 20 million Americans fall into this unsupervised category.

Our observation that the majority of users of unconventional therapy did not discuss this therapy with their medical doctors suggests a deficiency in current patient-doctor relations. Perhaps this lack of communication derives from medical doctors' mistaken assumption that their patients do not routinely use unconventional therapies for serious medical problems. Perhaps medical doctors do not discuss the use of unconventional therapies because they lack adequate knowledge of these techniques. In either case, this failure to communicate is not in the best interest of the patients, since the use of unconventional therapy, especially if it is totally unsupervised, may be harmful.<sup>11,27</sup>

Medical doctors should begin to ask their patients about their use of unconventional therapy whenever they obtain a history. Some doctors may be uncomfortable with this line of questioning. Nonetheless, as Kleinman and colleagues suggested more than a decade ago,<sup>28,29</sup> an exploration of the use of unconventional therapy and enhanced understanding of these practices will improve both communication between patients and doctors and clinical care. We suggest that medical schools include information about unconventional therapies and the clinical social sciences (anthropology and sociology) in their curriculums. The newly established National Institutes of Health Office for the Study of Unconventional Medical Practices should help promote scholarly research and education in this area.

We are indebted to Ms. Debi Arcaese for technical and editorial assistance.

#### REFERENCES

- Murray RH, Rubel AJ. Physicians and healers — unwitting partners in health care. *N Engl J Med* 1992;326:61-4.
- Gevitz N. Three perspectives on unorthodox medicine. In: Gevitz N, ed. *Other healers: unorthodox medicine in America*. Baltimore: Johns Hopkins University Press, 1988:1-28.
- Cleary PD. Chiropractic use: a test of several hypotheses. *Am J Public Health* 1982;72:727-30.
- McGuire MB. *Ritual healing in suburban America*. New Brunswick, N.J.: Rutgers University Press, 1988.
- Cook C, Baisden D. Ancillary use of folk medicine by patients in primary care clinics in southwestern West Virginia. *South Med J* 1986;79:1098-101.
- Cassileth BR, Lusk EJ, Strouse TB, Bodenheimer BJ. Contemporary unorthodox treatments in cancer medicine. *Ann Intern Med* 1984;101:105-12.
- U.S. Congress, Office of Technology Assessment. *Unconventional cancer treatments*. Washington, D.C.: Government Printing Office, 1990 (OTA-H-405).
- Clinical Oncology Group. New Zealand cancer patients and alternative medicine. *N Z Med J* 1987;100:110-3.
- Brigden ML. Unorthodox therapy and your cancer patient. *Postgrad Med* 1987;81:271-80.
- Cassileth BR. The social implications of questionable cancer therapies. *Cancer* 1989;63:1247-50.
- Harris and Associates. *Health, information and the use of questionable treatments: a study of the American public*. Washington, D.C.: Department of Health and Human Services, 1987.
- Cronan TA, Kaplan RM, Posner L, Blumberg E, Kozin F. Prevalence of the use of unconventional remedies for arthritis in a metropolitan community. *Arthritis Rheum* 1989;32:1604-7.
- Kronenfeld JJ, Wasner C. The use of unorthodox therapies and marginal practitioners. *Soc Sci Med* 1982;16:1119-25.
- Cherkin DC, MacCornack FA. Patient evaluations of low back pain care from family physicians and chiropractors. *West J Med* 1989;150:351-5.
- Cohen CJ, Eisenberg DM, Mayer KH, Delbanco TL. Prevalence of non-conventional medical treatments in HIV-infected patients: implications for primary care. *Clin Res* 1990;38:692A. abstract.
- Smart HL, Mayberry JF, Atkinson M. Alternative medicine consultations and remedies in patients with the irritable bowel syndrome. *Gut* 1986;27:826-8.
- Verhoef MJ, Sutherland LR, Brklich L. Use of alternative medicine by patients attending a gastroenterology clinic. *Can Med Assoc J* 1990;142:121-5.
- Snyder P. The use of nonprescribed treatments by hemodialysis patients. *Cult Med Psychiatry* 1983;7:57-76.
- Yager J, Landsverk J, Edelstein CK. Help seeking and satisfaction with care in 641 women with eating disorders. I. Patterns of utilization, attributed change, and perceived efficacy of treatment. *J Nerv Ment Dis* 1989;177:632-7.
- Gillick MR. Common-sense models of health and disease. *N Engl J Med* 1985;313:700-3.
- Lepkowski JM. Telephone sampling methods in the United States. In: Groves RM, Biemer PP, Lyberg LE, Massey JT, Nicholls WL II, Waksberg J, eds. *Telephone survey methodology*. New York: John Wiley, 1988:73-98.
- SUDAAN: professional software for survey data analysis, version 5.3. Research Triangle Park, N.C.: Research Triangle Institute, 1989.
- National health interview survey. Hyattsville, Md.: National Center for Health Statistics, 1989 (electronic data file).
- Schappert SM. National ambulatory medical care survey: 1990 summary. Advance data from vital and health statistics. No. 213. Hyattsville, Md.: National Center for Health Statistics, 1992. (DHHS publication no. (PHS) 92-1250.)
- Levit KR, Lazenby HC, Cowan CA, Letsch SW. National health expenditures, 1990. *Health Care Financ Rev* 1991;13(1):29-54.
- National Center for Health Statistics. Health and nutrition examination survey II, 1976-1980: medical history ages 12-74 years (ICPSR 8183). 2nd ICPSR ed. Ann Arbor, Mich.: Inter-university Consortium for Political and Social Research, 1984 (electronic data file).
- Subcommittee on Health and Long-Term Care of the Select Committee on Aging, House of Representatives, Ninety-Eighth Congress, Second Session. *Quackery: a \$10 billion scandal*. Washington, D.C.: Government Printing Office, 1984.
- Kleinman A, Eisenberg L, Good B. Culture, illness, and care: clinical lessons from anthropologic and cross-cultural research. *Ann Intern Med* 1978;88:251-8.
- Culture, health care systems, and clinical reality. In: Kleinman A. *Patients and healers in the context of culture: an exploration of the borderland between anthropology, medicine, and psychiatry*. Berkeley: University of California Press, 1980:50-60.