



ECZEMA

Eczema is probably one of the most difficult dermatological conditions to treat in a practical manner. The difficulty in treatment lies mostly in the fact that there are so many forms of eczema: acute, chronic, allergic, dry, weeping, impetiginous, seborrheic. The causes are sometimes elusive; many pollutants and environmental toxins contribute to the insidious development of eczema. In an attempt to provide several options in the treatment of this common complaint, antihomotoxic preparations commensurate to some of the different forms of eczema are discussed as well as possible protocols to use.

One of the main products used in eczema is Graphites-Homaccord, mainly because it can be used for a long term therapy. According to Dahlke, "throughout the whole remedy, there is a characteristic tendency towards the formation of cracks." Engystol is another basic preparation for eczema, particularly for infected or allergic eczema, as Engystol vanquishes many of the toxins of eczema.

The following chart illustrates several antihomotoxic preparations specific to the treatment of eczema. With these the practitioner can design a biological treatment plan specific to the individual case and its underlying cause.

TYPE OF ECZEMA

APPLICABLE ANTIHOMOTOXIC PREPARATIONS POSSIBLE PROTOCOL*

Acute

GRAPHITES-HOMACCORD and GALIUM-HEEL

1 ampule of each i.v.; s.c.; or orally 3 x per week for 3 weeks; then twice a week for 3 weeks, then once a week for at least 3 weeks.

Allergic

ENGYSTOL

Engystol i.v. 2 x per week for 2 weeks, then once a week for 5 weeks. Continue oral therapy with tablets or ampules: 3-5 tabs daily or one ampule/day for 5-8 weeks.

Food related

LYCOPodium-INJEEL, APIS-HOMACCORD, ENGYSTOL, HEPAR COMPOSITUM

First injection i.v. 1 amp. Lycopodium + 1 amp. Hepar comp then orally: 1 amp. Lycopodium + 1 amp. Engystol + 1 amp. Hepar comp 3 x week for 1 week, then 2 x week for 3 weeks.

Dry

GRAPHITES-HOMACCORD

Graphites-Homaccord drops: 10 drops 3 x a day for 5 weeks.

Of the palms (hands)

HEPAR COMPOSITUM

1 ampule Hepar comp i.v. once a week for 3 weeks + 1 ampule Hepar comp orally 2 x a week (in addition to injection) for 4 weeks.

Pustular

ACIDUM SULFURICUM INJEEL FORTE, CIMICIFUGA-HOMACCORD

1 ampule Acidum sulfuricum Injeel forte + 1 ampule Cimicifuga-Homaccord s.c. or orally once a week for 6-9 weeks.

Of the scalp

GRAPHITES-HOMACCORD, ACIDUM SULFURICUM INJEEL FORTE

10 drops Graphites-Homaccord + 1 ampule Acidum sulfuricum Injeel forte orally once a day for 10 weeks.

Scrotal or face

TRAUMEEL

1 ampule Traumeel i.v. or s.c. or orally 2 x week for 5 weeks.

Seborrheic

UBIQUINON COMPOSITUM, UBICOENZYME, NATRUM MURIATICUM INJEEL FORTE, OVARIVM COMPOSITUM

1 ampule Ubiquinon comp + 1 ampule Natrum muriaticum Injeel + 1 ampule Ovarium comp orally 2 x week for 8 weeks.

* All remedies can be given orally; injection therapy is aimed to give orthodox physicians the option of clinical administration.