

Traumeel S Gel

Empirical Report on its Use in Animals

Animal species	Possible Indications	Frequency and duration of treatment	Additional medication	Result	Tolerance	Adverse reactions	Assessment of efficacy
Cattle	Tarsitis (inflammation of the ankle joint) Carpalis (inflammation and swelling of the "knee" on the front leg) Phlegmon Haemaloma (bruise) Omphalitis (inflammation of the umbilicus) Mastitis Soft tissue swelling (e.g. after contusions, difficult births)	All animal species were treated topically 2 to 3 times daily for 1-2 weeks	As additional medication, Traumeel S was administered internally or per injection. If necessary, other homeopathic products (e.g. Arnica-Heel, Apis-Homaccord) or antibiotics were used.	After 1 or 2 weeks of treatment, the symptoms of inflammation or local symptoms subsided.	In all cases good, in most cases very good. Signs of intolerance were not observed.	none	In all cases good to very good
Horse	Tendinitis, tendovaginitis Thrombophlebitis (inflammation of a vein, usually the cervical vein) Bruise, contusion Pressure from saddle Carpalis, tarsitis						
Dog	Distortion (sprain of a joint) Arthritis Myositis (inflammation of a muscle, hardening) Haemaloma, seroma Myogenic cervical syndrome (muscular hardening in neck region)						
Cat	Arthritis Haemaloma, seroma						